



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Substance.	Mean refraction ($\mu_D - 1$).	Specific gravity.	Ratio.
Ice	0.3089	0.9184	2973
Water at 0° C.	0.3330	0.9993	3001

Moreover, as a general rule, those substances that are most affected in density by heat are the most sensitive.

6. No sudden change of sensitiveness occurs near the boiling-point; at least this is true in respect to bisulphide of carbon, ether, and methylic alcohol.

XVIII. "On the Adaptation of the Human Eye to varying Distances." By CHARLES ARCHER, Esq., Surgeon, Bengal Army. Communicated by Prof. STOKES, Sec. R.S.
Received June 17, 1858.

(Abstract.)

The following is a summary of the author's views on the question :—

1. The eye is adapted to varying distances principally by an alteration in the fibrous arrangement of the lens itself. Moreover, that when the lens is removed after an operation for cataract, the power of adaptation is nearly lost, and can only be exerted within very confined distances.

2. That the purpose of focalizing light at short distances is doubtless assisted, as suggested by Bowman, by the contractions of the ciliary muscle, in its antero-posterior direction, bringing forward the ciliary processes.

3. That as the posterior hemisphere of the capsule is firmly united to the hyaloid membrane, this portion must always remain quiescent, and therefore the antero-posterior contractions of the ciliary muscle must be very limited as regards the lens.

4. That the ciliary muscle, being placed around the eye, and its fibres being of a somewhat plexiform character, the contractions of the muscle will relax those yielding portions of the eye placed within its circumference.

5. That the relaxations of the ciliary processes will deprive the capsule of its firm support. It will be pressed forward by the lens,

which will meet with no further resistance to the expansion of its short axis.

6. That the lens itself, as microscopically described by Bowman and Kölliker, is admirably adapted to the varying changes which take place in the capsule.

7. That the posterior capsule being firmly united to the hyaloid membrane, the alteration in the diameters of the cavity of the capsule must take place from the periphery of the lens to its centre, and from behind forwards, but not from before backwards, on account of the close union of the posterior capsule to the hyaloid membrane.

8. That to allow such alteration to take place without endangering the achromatism of the lens, the alterations in the plane of its long diameter must be synchronous with the alterations in the plane of its short diameter. To allow of this, the margin of the lens is free in the canal of Petit; were it not the case, chromatic aberration would result.

9. That the elasticity of the capsule of the lens and the ciliary muscle are antagonistic; that on the ciliary muscle becoming relaxed, the capsule of the lens is free to exert that elasticity.

10. That, by the pressure exerted by the anterior hemisphere of the capsule by means of the polygonal cells of Virchow on the anterior face of the lens, the organ is able to fulfil all the requirements for adapting it to receive focalized light from long distances.

11. That the polygonal cells of Virchow are placed on the posterior surface of the anterior hemisphere of the capsule with the view before mentioned, and that they are arranged with their long diameters in an antero-posterior direction, that pressure may not injure their transparency, which would be the case if placed laterally.

12. That these cells are not found in other parts of the capsule.

13. That the fibres of the lens are serrated for the purpose of uniting either to other, so as to allow them greater freedom of motion without altering their ultimate relations to each other.

14. That the ciliary muscle is very highly endowed with nervous matter to supply all these varying requirements.

15. By the above postulates, all the modern discoveries in the microscopical anatomy of the eye receive a distinct expression of their individual functions, and, by so doing, adapt the organ of vision to the acknowledged laws of light.